

## UTILIZATION MANAGEMENT MEDICAL POLICY

**POLICY:** Hemophilia – Gene Therapy – Hemgenix Utilization Management Medical Policy

- Hemgenix® (etranacogene dezaparvovec-drlb intravenous infusion – CSL Behring and uniQure)

**REVIEW DATE:** 02/28/2024

---

### OVERVIEW

Hemgenix, an adeno-associated virus vector-based gene therapy, is indicated for the treatment of adults with hemophilia B (congenital Factor IX deficiency) who: 1) currently use Factor IX prophylaxis therapy; or 2) have current or historical life-threatening hemorrhage; or 3) have repeated, serious spontaneous bleeding episodes.<sup>1</sup> The recommended dose of Hemgenix is  $2 \times 10^{13}$  genome copies per kg of body weight given as an one-time (per lifetime) single dose as an intravenous infusion.

### Disease Overview

Hemophilia B is a genetic bleeding disorder caused by missing or insufficient levels of blood Factor IX, a protein required to produce blood clots to halt bleeding.<sup>2-5</sup> The condition is a rare X-linked bleeding disorder that mainly impacts males. Hemophilia B is four times less common than hemophilia A, which is caused by a relative lack of blood Factor VIII. Approximately 30,000 individuals are living with hemophilia in the US and hemophilia B accounts for around 15% to 20% of hemophilia cases, or around 6,000 patients. Symptoms patients may experience include heavy or prolonged bleeding following an injury or after a medical procedure. Bleeding can also occur internally into joints, muscles, or internal organs. Spontaneous bleeding events may also occur. Complications in patients with hemophilia B include joint disease and hemarthrosis. Hemophilia B may be diagnosed when bleeding occurs in infancy or later in life for those with milder disease. There is a strong correlation between Factor IX levels and phenotypic expression of bleeding. Normal plasma levels of Factor IX range from 50% to 150%. The disease is classified based on reduced levels. Mild, moderate, and severe hemophilia B are characterized by Factor IX levels ranging from 6% up to 49%, 1% up to 5%, and < 1%, respectively. Besides Hemgenix, Factor IX products, both recombinant and plasma-derived, are used routinely to prevent bleeding or are given on demand to treat bleeding episodes associated with hemophilia B.

### Clinical Efficacy

The efficacy of Hemgenix was evaluated in a prospective, open-label, single-dose, single-arm, multinational pivotal study called HOPE-B that involved 54 adult male patients with moderately severe or severe hemophilia B (Factor IX levels  $\leq 2\%$ ).<sup>1</sup> Patients prospectively completed a lead-in period of at least 6 months in which standard care routine Factor IX prophylaxis therapy was given. This was followed by a single intravenous dose of  $2 \times 10^{13}$  genome copies/kg of body weight of Hemgenix. Patients were permitted to continue Factor IX prophylaxis during Months 0 to 6 after dosing, if needed, until Factor IX levels were adequate. Prior to screening, patients had been on stable prophylactic therapy for at least 2 months and had greater than 150 exposure days of treatment with a Factor IX product.<sup>6</sup> Factor IX inhibitors (or a history), uncontrolled HIV, or advanced liver fibrosis prevented participation. Adequate hepatic and renal function were required. The estimated mean annualized bleeding rate during Months 7 to 18 following Hemgenix treatment was 1.9 bleeds/year compared with 4.1 bleeds/year during the lead-in period (before Hemgenix administration).<sup>1</sup> At 18 months after treatment, Factor IX activity had increased by 34.3%. The HOPE-B trial is ongoing.

### POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Hemgenix. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indication. Because of the specialized skills required for evaluation and diagnosis of patients treated with Hemgenix as well as the monitoring required for adverse events and long-term efficacy, approval requires Hemgenix to be prescribed by a physician who specializes in the condition being treated. All approvals are provided for one-time (per lifetime) as a single dose. If claims history is available, verification is required for certain criteria as noted by **[verification in claims history required]**. For the dosing criteria, verification of the appropriate weight-based dosing is required by a Medical Director as noted by **[verification required]**. In the criteria for Hemgenix, as appropriate, an asterisk (\*) is noted next to the specified gender. In this context, the specified gender is defined as follows: males are defined as individuals with the biological traits of a man, regardless of the individual's gender identity or gender expression. All reviews (approvals and denials) will be forwarded to the Medical Director for evaluation.

Some clients have elected Embarc Benefit Protection. For these clients, the Medical Director will coordinate with eviCore to ensure the Embarc Benefit Protection portion of the review has been completed. If the Embarc Benefit Protection portion of the review has not been completed, the Medical Director will route to [Embarc@eviCore.com](mailto:Embarc@eviCore.com) prior to completing the review.

**Documentation:** Documentation is required for use of Hemgenix as noted in the criteria as **[documentation required]**. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information.

**Automation:** None.

## RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Hemgenix is recommended in those who meet the following criteria:

### FDA-Approved Indication

- 
1. **Hemophilia B.** Approve a one-time (per lifetime) single dose if the patient meets ALL of the following (A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, and Q):
    - A) Patient is male\*; AND
    - B) Patient is  $\geq 18$  years of age; AND
    - C) Patient has not received Hemgenix in the past **[verification in claims history required]**; AND  
Note: If no claim for Hemgenix is present (or if claims history is not available), the prescribing physician confirms that the patient has not previously received Hemgenix.
    - D) Patient has moderately severe or severe hemophilia B as evidence by a baseline (without Factor IX replacement therapy) Factor IX level of  $\leq 2\%$  of normal **[documentation required]**; AND
    - E) Patient meets ONE of the following (i, ii, or iii):
      - i. Patient meets BOTH of the following (a and b):
        - a) Patient has been receiving routine prophylaxis with Factor IX therapy continuously for  $\geq 2$  months **[documentation required]**; AND
        - b) According to the prescribing physician, the patient has a history of use of Factor IX therapy for  $\geq 150$  exposure days; OR
      - ii. Patient meets BOTH of the following (a and b):
        - a) Patient has a history of life-threatening hemorrhage; AND
        - b) On-demand use of Factor IX therapy was required for this life-threatening hemorrhage; OR
      - iii. Patient meets BOTH of the following (a and b):

- a) Patient has a history of repeated, serious spontaneous bleeding episodes; AND
- b) On-demand use of Factor IX therapy was required for these serious spontaneous bleeding episodes; AND
- F) Patient meets ALL of the following (i, ii, and iii):
  - i. Factor IX inhibitor titer testing has been performed within 30 days **[documentation required]**; AND
  - ii. Patient does not currently have an inhibitor to Factor IX **[documentation required]**; AND
  - iii. Patient does not have a history of Factor IX inhibitors **[documentation required]**; AND
- G) Prophylactic therapy with Factor IX will not be given after Hemgenix administration once adequate Factor IX levels have been achieved; AND  
Note: Use of episodic Factor IX therapy is acceptable for the treatment of bleeds and for surgery/procedures if needed as determined by the hemophilia specialist physician.
- H) Patient meets BOTH of the following (i and ii):
  - i. Patient does not have an active infection with hepatitis B virus or hepatitis C virus **[documentation required]**; AND
  - ii. Patient is not currently receiving antiviral therapy for a prior hepatitis B virus or hepatitis C virus exposure **[documentation required]**; AND
- I) Patient does not have uncontrolled human immunodeficiency virus infection **[documentation required]**; AND  
Note: A patient testing positive for human immunodeficiency virus can still qualify for Hemgenix if controlled on antiviral therapy with CD4+ counts  $\geq 200/\mu\text{L}$  or a viral load  $\leq 200$  copies/mL.
- J) Patient has undergone liver function testing within 30 days and meets ALL of the following (i, ii, and iv):
  - i. Alanine aminotransferase level is  $\leq$  two times the upper limit of normal **[documentation required]**; AND
  - ii. Aspartate aminotransferase level is  $\leq$  two times the upper limit of normal **[documentation required]**; AND
  - iii. Total bilirubin level is  $\leq$  two times the upper limit of normal **[documentation required]**; AND
  - iv. Alkaline phosphatase level is  $\leq$  two times the upper limit of normal **[documentation required]**; AND
- K) Patient does not have evidence of advanced liver impairment and/or advanced fibrosis; AND  
Note: For example, liver elastography (e.g.,  $\geq 9$  kPa) suggestive of or equal to METAVIR Stage 3 disease.
- L) Within 30 days, the platelet count was  $\geq 50 \times 10^9/\text{L}$  **[documentation required]**; AND
- M) Within 30 days, patient meets BOTH of the following (i and ii):
  - i. Patient has an estimated creatinine clearance  $\geq 30$  mL/min **[documentation required]**; AND
  - ii. Creatinine level is  $\leq$  two times the upper limit of normal **[documentation required]**; AND
- N) Patient does not have another coagulation disorder, besides hemophilia B; AND
- O) The medication is prescribed by a hemophilia specialist physician; AND
- P) Current patient body weight has been obtained within 30 days **[documentation required]**; AND
- Q) If criteria A through P are met, approve one dose (kit) of Hemgenix to provide for a one-time (per lifetime) single dose of  $2 \times 10^{13}$  genome copies per kg of body weight by intravenous infusion **[verification required]**. Table 1 provides the kit size and the National Drug Codes (NDCs).

\* Refer to the Policy Statement.

**Dosing.** The recommended dose of Hemgenix is a one-time (per lifetime) single dose of  $2 \times 10^{13}$  genome copies per kg of body weight by intravenous infusion.

**CONDITIONS NOT RECOMMENDED FOR APPROVAL**

Coverage of Hemgenix is not recommended in the following situations:

- 1. Prior Receipt of Gene Therapy.** Prior receipt of gene therapy was a reason for patient exclusion in the pivotal study.
- 2.** Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

**Table 1. Hemgenix Multi-Vial Kits.<sup>1</sup>**

Total Number of Vials per Kit	Patient Body Weight	Total Volume per Kit	NDC Number
10	46 to 50 kg	100	0053-0100-10
11	51 to 55 kg	110	0053-0110-11
12	56 to 60 kg	120	0053-0120-12
13	61 to 65 kg	130	0053-0130-13
14	66 to 70 kg	140	0053-0140-14
15	71 to 75 kg	150	0053-0150-15
16	76 to 80 kg	160	0053-0160-16
17	81 to 85 kg	170	0053-0170-17
18	86 to 90 kg	180	0053-0180-18
19	91 to 95 kg	190	0053-0190-19
20	96 to 100 kg	200	0053-0200-20
21	101 to 105 kg	210	0053-0210-21
22	106 to 110 kg	220	0053-0220-22
23	111 to 115 kg	230	0053-0230-23
24	116 to 120 kg	240	0053-0240-24
25	121 to 125 kg	250	0053-0250-25
26	126 to 130 kg	260	0053-0260-26
27	131 to 135 kg	270	0053-0270-27
28	136 to 140 kg	280	0053-0280-28
29	141 to 145 kg	290	0053-0290-29
30	146 to 150 kg	300	0053-0300-30
31	151 to 155 kg	310	0053-0310-31
32	156 to 160 kg	320	0053-0320-32
33	161 to 165 kg	330	0053-0330-33
34	166 to 170 kg	340	0053-0340-34
35	171 to 175 kg	350	0053-0350-35
36	176 to 180 kg	360	0053-0360-36
37	181 to 185 kg	370	0053-0370-37
38	186 to 190 kg	380	0053-0380-38
39	191 to 195 kg	390	0053-0390-39
40	196 to 200 kg	400	0053-0400-40
41	201 to 205 kg	410	0053-0410-41
42	206 to 210 kg	420	0053-0420-42
43	211 to 215 kg	430	0053-0430-43
44	216 to 220 kg	440	0053-0440-44
45	221 to 225 kg	450	0053-0450-45
46	226 to 230 kg	460	0053-0460-46
47	231 to 235 kg	470	0053-0470-47
48	236 to 240 kg	480	0053-0480-48

NDC – National Drug Code.

## REFERENCES

1. Hemgenix® intravenous infusion [prescribing information]. King of Prussia, PA; Kankakee, IL; and Lexington, MA: CSL Behring and uniQure; November 2022.
2. National Bleeding Disorders Foundation. Hemophilia B. An overview of symptoms, genetics, and treatments to help you understand hemophilia B. Available at: <https://www.hemophilia.org/bleeding-disorders-a-z/types/hemophilia-b>. Accessed on February 28, 2024.
3. Sidonio RF, Malec L. Hemophilia (Factor IX deficiency). *Hematol Oncol Clin N Am*. 2021;35:1143-1155.
4. Mancuso ME, Mahlangu JN, Pipe SW. The changing treatment landscape in haemophilia: from standard half-life clotting factor concentrates to gene editing. *Lancet*. 2021;397:630-640.
5. Croteau SE. Hemophilia A/B. *Hematol Oncol Clin N Am*. 2022;36:797-812.
6. Pipe SW, Leebeek FWG, Recht M, et al. Gene therapy with etranacogene dexaparvovec for hemophilia B. *N Engl J Med*. 2023;388:706-718.

## HISTORY

Type of Revision	Summary of Changes	Review Date
New Policy	--	01/11/2023
Annual Revision	<p>The following changes were made:</p> <p><b>Hemophilia B:</b> An overview of the changes are described below.</p> <ul style="list-style-type: none"> <li>• The documentation requirement was removed regarding the criterion that the “Patient does not have evidence of advanced liver impairment and/or advanced fibrosis”.</li> <li>• The following criteria were removed which stated that after the Hemgenix infusion, the physician attests that the following will be performed: 1) liver enzyme testing to monitor for liver enzyme elevations will be done at least weekly for the first 3 months and periodically thereafter; AND implementing a course of corticosteroids will be considered if the patient experiences clinically relevant increases in alanine aminotransferase levels; 2) the patient will undergo monitoring for Factor IX activity at least weekly for the first 3 months and periodically thereafter; and 3) the patient with preexisting risk factors for hepatocellular carcinoma will receive abdominal ultrasound screenings and be monitored at least annually for alpha fetoprotein elevations in the 5 years following receipt of Hemgenix.</li> <li>• The requirement for the specialist physician was changed from “physician who specializes in hemophilia” to “hemophilia specialist physician”.</li> <li>• The criterion regarding a current patient body weight be obtained within 30 days was moved to a separate criterion. Previously, this requirement was combined with the Dosing.</li> <li>• Dosing was clarified with emphasis that Hemgenix is given as a “single dose”. Also, “documentation required” was replaced with “verification required”. A related sentence was added to the Policy Statement that verification of the appropriate weight-based dosing is required by the Medical Director.</li> <li>• Regarding use of Hemgenix in the past, the phrase “verification required by prescriber” was changed to “verification in claims history required”. A qualifier was added to reflect that this requirement applies only if a claims history is available; this change was also reflected in the related Policy Statement. Wording regarding “prescriber must attest” was changed to “prescribing physician confirms” regarding the verification that the patient has not previously received Hemgenix. Also, in the Note, the following statement was removed as it is duplicative: verify through claims history that the patient has not previously received Hemgenix.</li> <li>• The phrase “prescriber attests” was removed from the requirement that “prophylactic therapy with Factor IX will not be given after Hemgenix administration once adequate Factor IX levels have been achieved” as well as the to the requirement regarding “patient does not have another coagulation disorder, besides hemophilia B”.</li> <li>• In the requirement that Factor IX inhibitor titer testing has been performed “within 30 days”, the phrase “before receipt of Hemgenix” was removed.</li> <li>• The phrase regarding liver “health assessment” was changed to liver “function testing”.</li> <li>• For the requirement that the patient does not have uncontrolled human immunodeficiency virus, the word “infection” was added after this phrase.</li> </ul> <p><b>Conditions Not Recommended for Approval:</b> The condition of “Prior Receipt of Gene Therapy” was added.</p>	02/28/2024